



# Recombinant Human Eukaryotic translation initiation factor 6 (EIF6)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-YP007582HU  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P56537  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Homo sapiens (Human)  |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | MAVRASFENN CEIGCFAKLT NTYCLVAIGG SENFYSVFEG ELSDTIPVVH<br>ASIAGCRIIG RMCVGNRHGL LVPNNTTDE LQHIRNSLPD TVQIRRVEER<br>LSALGNVTTC NDYVALVHPD LDRETEEILA DVLKVEVFRQ TVADQVLVGS<br>YCVFSNQGGL VHPKTSIEDQ DELSSLLQVP LVAGTVNRGS EVIAAGMVVN<br>DWCAFCGLDT TSTELSVVES VFKLNEAQPS TIATSMRDSL IDSLT  |
| <b>Source</b>            | Yeast   |
| <b>Target Names</b>      | EIF6  |
| <b>Protein Names</b>     | Recommended name: Eukaryotic translation initiation factor 6 Short name=<br>eIF-6 Alternative name(s): B(2)GCN homolog B4 integrin interactor CAB<br>p27(BBP)   |
| <b>Expression Region</b> | 1-245   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at<br>4°C for up to one week.  |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Full length protein   |
| <b>Target Details</b>    | Hemidesmosomes are structures which link the basal lamina to the intermediate<br>filament cytoskeleton. An important functional component of hemidesmosomes<br>is the integrin beta-4 subunit (ITGB4), a protein containing two fibronectin type<br>III domains. This protein binds to the fibronectin type III domains of ITGB4 and<br>may help link ITGB4 to the intermediate filament cytoskeleton. The encoded<br>protein, which is insoluble and found both in the nucleus and in the cytoplasm,<br>can function as a translation initiation factor and prevent the association of the<br>40S and 60S ribosomal subunits. Multiple transcript variants encoding two<br>different isoforms have been found for this gene. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the<br>contents to the bottom. Please reconstitute protein in deionized sterile water to a<br>concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final<br>concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final<br>concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients,<br>storage temperature and the stability of the protein itself.   |



Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.