




CAPN2 Antibody

Product Code	CSB-PA004496LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P17655
Immunogen	Recombinant Human Calpain-2 catalytic subunit protein (20-700AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF, IP; Recommended dilution: WB:1:1000-1:5000, IHC:1:200-1:500, IF:1:50-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Calpain-2 catalytic subunit (EC 3.4.22.53) (Calcium-activated neutral proteinase 2) (CANP 2) (Calpain M-type) (Calpain large polypeptide L2) (Calpain-2 large subunit) (Millimolar-calpain) (M-calpain), CAPN2, CANPL2
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	CAPN2

Image



Western Blot

Positive WB detected in: 293T whole cell lysate


All lanes: CAPN2 antibody at 3.4µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 80, 72 kDa


Observed band size: 80 kDa




IHC image of CSB-PA004496LA01HU diluted at 1:300 and staining in paraffin-embedded human tonsil tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA004496LA01HU diluted at 1:300 and staining in paraffin-embedded human appendix tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HepG2 cells with CSB-PA004496LA01HU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunoprecipitating CAPN2 in Hela whole cell lysate
 Lane 1: Rabbit control IgG instead of CSB-PA004496LA01HU in Hela whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)
 Lane 2: CSB-PA004496LA01HU (8μg) + Hela whole cell lysate (500μg)
 Lane 3: Hela whole cell lysate (10μg)

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.