




Rabbit anti-Human IgG Fc Antibody;HRP conjugated

Product Code	CSB-PA00540F0Rb
Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze.
Immunogen	Human IgG Fc fragment
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB; Recommended dilution: WB:1:500-1:5000
Relevance	The fragment crystallizable region (Fc region) is the tail region of an antibody that interacts with cell surface receptors called Fc receptors and some proteins of the complement system. This property allows antibodies to activate the immune system. In IgG, IgA and IgD antibody isotypes, the Fc region is composed of two identical protein fragments, derived from the second and third constant domains of the antibody two heavy chains; IgM and IgE Fc regions contain three heavy chain constant domains (CH domains 2–4) in each polypeptide chain. The Fc regions of IgGs bear a highly conserved N-glycosylation site. Glycosylation of the Fc fragment is essential for Fc receptor-mediated activity. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and α -2,6 linked sialic acid residues.
Form	Liquid
Conjugate	HRP conjugated
Storage Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Purification Method	Caprylic Acid Ammonium Sulfate Precipitation purified
Isotype	IgG
Clonality	Polyclonal
Product Type	Secondary Antibody

Image



Western Blot

Positive WB detected in Recombinant protein (80ng, 40ng, 20ng, 10ng)

All lanes: Human IgG Fc antibody; HRP conjugated at 1:1000

Predicted band size: 35 kDa

Observed band size: 35 kDa



Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.