



Rabbit anti-Human IgG Fc Antibody; Biotin conjugated

Product Code	CSB-PA00540H0Rb
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Immunogen	Human IgG Fc fragment
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	<p>The fragment crystallizable region (Fc region) is the tail region of an antibody that interacts with cell surface receptors called Fc receptors and some proteins of the complement system. This property allows antibodies to activate the immune system. In IgG, IgA and IgD antibody isotypes, the Fc region is composed of two identical protein fragments, derived from the second and third constant domains of the antibody two heavy chains; IgM and IgE Fc regions contain three heavy chain constant domains (CH domains 2–4) in each polypeptide chain. The Fc regions of IgGs bear a highly conserved N-glycosylation site. Glycosylation of the Fc fragment is essential for Fc receptor-mediated activity. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and α-2,6 linked sialic acid residues.</p>
Form	Liquid
Conjugate	Biotin conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Caprylic Acid Ammonium Sulfate Precipitation purified
Isotype	IgG
Clonality	Polyclonal
Alias	fragment crystallizable region
Product Type	Secondary Antibody