




VEGFD Antibody

| | |
|----------------------------|---|
| Product Code | CSB-PA07554A0Rb |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | O43915 |
| Immunogen | Recombinant Human Vascular endothelial growth factor D protein (89-205AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC; Recommended dilution: IHC:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.1mol/L NaCl, 15mmol/L NaN ₃ , PH7.2 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | c-fos induced growth factor antibody; c-fos induced growth factor (vascular endothelial growth factor D) antibody; c-fos induced growth factor antibody; c-fos-induced growth factor antibody; FIGF antibody; Vascular endothelial growth factor D antibody; Vascular endothelial growth factor D precursor antibody; Vascular endothelial growth factor D precursor antibody; VEGF D antibody; VEGF-D antibody; VEGFD antibody; VEGFD_HUMAN antibody |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Cardiovascular |
| Target Names | VEGFD |

Image



IHC image of CSB-PA07554A0Rb diluted at 1:225 and staining in paraffin-embedded human heart tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.