



# Recombinant Dog Cytochrome c oxidase subunit 6A2, mitochondrial (COX6A2)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-MP005849DO  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P61900  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Canis lupus familiaris (Dog) (Canis familiaris)   |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | ASGAKGDHGG AGASTXXLLT   |
| <b>Source</b>            | Mammalian cell  |
| <b>Target Names</b>      | COX6A2  |
| <b>Protein Names</b>     | Recommended name: Cytochrome c oxidase subunit 6A2, mitochondrial<br>Alternative name(s): Cytochrome c oxidase polypeptide VIa-heart Short name= COXVIAH  |
| <b>Expression Region</b> | 1-20  |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Full length protein   |
| <b>Target Details</b>    | Cytochrome c oxidase (COX), the terminal enzyme of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may be involved in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (heart/muscle isoform) of subunit VIa, and polypeptide 2 is present only in striated muscles. Polypeptide 1 (liver isoform) of subunit VIa is encoded by a different gene, and is found in all non-muscle tissues. These two polypeptides share 66% amino acid sequence identity. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life  |



of lyophilized form is 12 months at -20°C/-80°C.