



# Recombinant Dog Cell division control protein 42 homolog (CDC42)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-EP005008DO  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P60952  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Canis lupus familiaris (Dog) (Canis familiaris)   |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | MQTIKCVVVG DGAVGKTCLL ISYTTNKFPS EYVPTVFDNY AVTVMIGGEP<br>YTLGLFDTAG QEDYDRLRPL SYPQTDVFLV CFSVVSPSSF ENVKEKWWPE<br>ITHHCPKTPF LLVGTQIDLR DDPSTIEKLA KNKQKPITPE TAEKLARDLK<br>AVKYVECSAL TQKGLKNVFD EAILAALEPP EPKKSRRC   |
| <b>Source</b>            | E.coli  |
| <b>Target Names</b>      | CDC42   |
| <b>Protein Names</b>     | Recommended name: Cell division control protein 42 homolog<br>Alternative name(s): G25K GTP-binding protein   |
| <b>Expression Region</b> | 1-188   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Full length protein   |
| <b>Target Details</b>    | This protein is a small GTPase of the Rho-subfamily, which regulates signaling pathways that control diverse cellular functions including cell morphology, migration, endocytosis and cell cycle progression. This protein is highly similar to <i>Saccharomyces cerevisiae</i> Cdc 42, and is able to complement the yeast <i>cdc42-1</i> mutant. The product of oncogene <i>Dbl</i> was reported to specifically catalyze the dissociation of GDP from this protein. This protein could regulate actin polymerization through its direct binding to Neural Wiskott-Aldrich syndrome protein (N-WASP), which subsequently activates Arp2/3 complex. Alternative splicing of this gene results in multiple transcript variants. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life  |



of lyophilized form is 12 months at -20°C/-80°C.