



# Recombinant Human Disabled homolog 2 (DAB2)

<b>Product Code</b>	CSB-EP006480HU
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P98082
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	<p>SNEVETSAT NGQPDAQAAP KAPSKKEKKK GPEKTDEYLL ARFKGDGVKY  KAKLIGIDDV PDARGDKMSQ DSMMLKKGMA AAGRSQGQHK QRIWVNISLS  GIKIIDEKTG VIEHEHPVNK ISFIARDVTD NRAFGYVCGG EGQHQQFFAIK  TGQQAELVV DLKDLFQVIY NVKKKEEEKK KIEEASKAVE NGSEALMILD  DQTNKLKSGV DQMDLFGDMS TPPDLNSPTE SKDILLVDLN SEIDTNQNSL  RENPF LTNGI TSCSLPRPTP QASFLPENAF SANLNFFPTP NPD PFRDDPF  TQPDQSTPSS FDSLKSPDQK KENSSSSSTP LSNGLNGDV DYFGQQFDQI  SNRTGKQEAQ AGPWPFFSSQ TQPAVRTQNG VSEREQNGFS  VKSSPNPFVG SPPKGLSIQN GVKQDLESSV QSSPHDSIAI IPPPQSTKPG  RGRRTAKSSA NDLLASDIFA PPVSEPSGQA SPTGQPTALQ PNPLDLFKTS  APAPVGPLVG LGGVTVTL PQ AGPWNTASLV FNQSPSMAPG  AMMGGQPSGF SQPVIFGTSP AVSGWNQPSP FFASTPPPVP  VWGPSASVA PNAWSTTSPL GNPFQSNIFP APAVSTQPPS MHSSLLVTPP  QPPPRAGPPK DISSDAFTAL DPLGDKEIKD VKEMFKDFQL RQPPAVPARK  GEQTSSGTL S AFASYFNSKV GIPQENADHD DFDANQLLNK INEPPKPAPR  QVSLPVTKST DNAFENPFFK DSGFSSQASV ASSQPVSSEM YRDPFGNPFA</p>
<b>Source</b>	E.coli
<b>Target Names</b>	DAB2
<b>Protein Names</b>	Recommended name: Disabled homolog 2 Alternative name(s): DOC-2 Differentially-expressed protein 2
<b>Expression Region</b>	2-770
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	DAB2 mRNA is expressed in normal ovarian epithelial cells but is down-regulated or absent from ovarian carcinoma cell lines. The 770-amino acid predicted protein has an overall 83% identity with the mouse p96 protein, a putative mitogen-responsive phosphoprotein; homology is strongest in the amino-terminal end of the protein in a region corresponding to the phosphotyrosine interaction domain. The down-regulation of DAB2 may play an important role in ovarian carcinogenesis. This gene was initially named DOC2 (for Differentially expressed in Ovarian Cancer) and is distinct from the DOC2A and DOC2B genes (for double C2-like domains, alpha and beta).



---

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.