



# Recombinant Human Tyrosine 3-monoxygenase (TH)

|                          |  |
|--------------------------|--|
| <b>Product Code</b>      | CSB-BP023470HU   |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.  |
| <b>Uniprot No.</b>       | P07101   |
| <b>Product Type</b>      | Recombinant Protein  |
| <b>Immunogen Species</b> | Homo sapiens (Human)   |
| <b>Purity</b>            | ≥85% (SDS-PAGE)  |
| <b>Sequence</b>          | PTPDATTPQ AKGFRRVAVSE LDAQQAEAIM VRGQGAPGPS LTGSPWPGTA<br>APAASYTPTP RSPRFIGRRQ SLIEDARKER EAAVAAAAAA VPSEPGDPLE<br>AVAFEEKEGK AVLNLLFSPR ATKPSALSRA VKVFETFEAK IHHLETRPAQ<br>RPRAGGPHLE YFVRLEVRRG DLAALLSGVR QVSEDEVSPA GPKVPWFPRK<br>VSELDKCHHL VTKFDPDL DL DHPGFSDQVY RQRRKLIAEI AFQYRHGDPI<br>PRVEYTAEEI ATWKEVYTTL KGLYATHACG EHLEAFALLE RFSGYREDNI<br>PQLEDVSRFL KERTGFQLRP VAGLLSARDF LASLAFRVFQ CTQYIRHASS<br>PMHSPEPDCC HELLGHVPML ADRTFAQFSQ DIGLASLGAS DEEIEKLSTL<br>YWFTVEFGLC KQNGEVKAYG AGLLSSYGEL LHCLSEEPEI RAFDPEAAAV<br>QPYQDQTYQS VYFVSESFSD AKDKLRSYAS RIQRPFVSVKF DPYTLAIDVL<br>DSPQAVRRSL EGVQDELDTL AHALSAIG |
| <b>Source</b>            | Baculovirus  |
| <b>Target Names</b>      | TH   |
| <b>Protein Names</b>     | Recommended name: Tyrosine 3-monoxygenase EC= 1.14.16.2 Alternative name(s): Tyrosine 3-hydroxylase Short name= TH   |
| <b>Expression Region</b> | 2-528  |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.  |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.  |
| <b>Protein Length</b>    | Full Length of Mature Protein  |
| <b>Target Details</b>    | This protein is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in this gene have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.  |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.  |



## Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.