



# Recombinant Rabbit Myosin-7 (MYH7), partial

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-BP015300RB  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P04461  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Oryctolagus cuniculus (Rabbit)  |
| <b>Purity</b>            | ≥85% (SDS-PAGE)   |
| <b>Source</b>            | Baculovirus   |
| <b>Target Names</b>      | MYH7  |
| <b>Protein Names</b>     | Recommended name: Myosin-7 Alternative name(s): Beta isomyosin Myosin heavy chain 7 Myosin heavy chain slow isoform Short name= MyHC-slow Myosin heavy chain, cardiac muscle beta isoform Short name= MyHC-beta   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Partial   |
| <b>Target Details</b>    | Muscle myosin is a hexameric protein containing 2 heavy chain subunits, 2 alkali light chain subunits, and 2 regulatory light chain subunits. This gene encodes the beta (or slow) heavy chain subunit of cardiac myosin. It is expressed predominantly in normal human ventricle. It is also expressed in skeletal muscle tissues rich in slow-twitch type I muscle fibers. Changes in the relative abundance of this protein and the alpha (or fast) heavy subunit of cardiac myosin correlate with the contractile velocity of cardiac muscle. Its expression is also altered during thyroid hormone depletion and hemodynamic overloading. Mutations in this gene are associated with familial hypertrophic cardiomyopathy, myosin storage myopathy, dilated cardiomyopathy, and Laing early-onset distal myopathy. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.   |