



Recombinant Chicken Troponin I, fast skeletal muscle (TNNI2)

| | |
|--------------------------|---|
| Product Code | CSB-EP024012CH |
| Storage | Store at -20°C, for extended storage, conserve at -20°C or -80°C. |
| Uniprot No. | P68246 |
| Product Type | Recombinant Protein |
| Immunogen Species | Gallus gallus (Chicken) |
| Purity | >85% (SDS-PAGE) |
| Sequence | SDEEKRRRAATARRQHLSAMLQLAVTEIEKEAAAKEVEKQNYLAEHCPPLSL PGSMQEL QELCKKLHAKIDSVDEERYDTEVKLQKTNKELEDLSQKLFDLRGKFKRPPLRR VRMSADA MLRALLGSKHKVNMDLRANLKQVKKEDTEKEKDLRDVGDWRKNIEEKSGME GRKKMFEAG ES |
| Source | E.coli |
| Target Names | TNNI2 |
| Protein Names | Recommended name: Troponin I, fast skeletal muscle Alternative name(s): Troponin I, fast-twitch isoform |
| Expression Region | 2-183 |
| Notes | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| Tag Info | Tag type will be determined during the manufacturing process. |
| Protein Length | Full Length of Mature Protein |
| Target Details | This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final |



concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.