



Recombinant Bovine Cystatin-C (CST3)

Product Code	CSB-BP006091BO
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P01035
Product Type	Recombinant Protein
Immunogen Species	Bos taurus (Bovine)
Purity	≥85% (SDS-PAGE)
Sequence	QGPRKGRLLG GLMEADVNEE GVQEALSFV SEFNKRSNDA YQSRVVRVVR ARKQVVSGMN YFLDVELGRT TCTKSQANLD SCPFHNQPHL KREKLCSFQV YVVPWMNTIN LVKFSCQD
Source	Baculovirus
Target Names	CST3
Protein Names	Recommended name: Cystatin-C Alternative name(s): Colostrum thiol proteinase inhibitor Cystatin-3
Expression Region	31-148
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	<p>The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes the most abundant extracellular inhibitor of cysteine proteases, which is found in high concentrations in biological fluids and is expressed in virtually all organs of the body. A mutation in this gene has been associated with amyloid angiopathy. Expression of this protein in vascular wall smooth muscle cells is severely reduced in both atherosclerotic and aneurysmal aortic lesions, establishing its role in vascular disease.</p>
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.



Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.