



# Recombinant Human Cyclin-dependent kinase 2-associated protein 1 (CDK2AP1)

|                          |  |
|--------------------------|--|
| <b>Product Code</b>      | CSB-YP005062HU   |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.  |
| <b>Uniprot No.</b>       | O14519   |
| <b>Product Type</b>      | Recombinant Protein  |
| <b>Immunogen Species</b> | Homo sapiens (Human)   |
| <b>Purity</b>            | ≥85% (SDS-PAGE)  |
| <b>Sequence</b>          | MSYKPNLAAH MPAAALNAAG SVHSPSTSMA TSSQYRQLLS DYGPPSLGYT<br>QGTGNSQVPQ SKYAELLAI EELGKEIRPT YAGSKSAMER LKRGIIHARG<br>LVRECLAETE RNARS  |
| <b>Source</b>            | Yeast  |
| <b>Target Names</b>      | CDK2AP1  |
| <b>Protein Names</b>     | Recommended name: Cyclin-dependent kinase 2-associated protein 1 Short name= CDK2-associated protein 1 Alternative name(s): Deleted in oral cancer 1 Short name= DOC-1 Putative oral cancer suppressor   |
| <b>Expression Region</b> | 1-115  |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.  |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.  |
| <b>Protein Length</b>    | Full length protein  |
| <b>Target Details</b>    | This protein is a specific CDK2-associated protein, which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggested the regulatory role in DNA replication during S phase of the cell cycle. A similar gene in hamster was isolated from, and functions as a growth suppressor of normal keratinocytes. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.  |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.  |