



# Recombinant Mouse Mitogen-activated protein kinase kinase kinase 5 (Map3k5), partial

|                          |  |
|--------------------------|--|
| <b>Product Code</b>      | CSB-EP013427MO   |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.  |
| <b>Uniprot No.</b>       | O35099   |
| <b>Product Type</b>      | Recombinant Protein  |
| <b>Immunogen Species</b> | Mus musculus (Mouse)   |
| <b>Purity</b>            | >85% (SDS-PAGE)  |
| <b>Source</b>            | E.coli   |
| <b>Target Names</b>      | Map3k5   |
| <b>Protein Names</b>     | Recommended name: Mitogen-activated protein kinase kinase kinase 5 EC= 2.7.11.25 Alternative name(s): Apoptosis signal-regulating kinase 1 Short name= ASK-1 MAPK/ERK kinase kinase 5 Short name= MEK kinase 5 Short name=   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.  |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.  |
| <b>Protein Length</b>    | Partial  |
| <b>Target Details</b>    | Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.  |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life   |



of lyophilized form is 12 months at -20°C/-80°C.