



# Recombinant Mouse Calpain-2 catalytic subunit (Capn2)

<b>Product Code</b>	CSB-YP004496MO
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	O08529
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Mus musculus (Mouse)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	S HERAIKYLNQ DYETLRNECL EAGALFQDPS FPALPSSSLGY KELGPYSSKT RGIEWKRPT E ICADPQFIIG GATRTRDQCQ ALGDCWLLAA IASLTLNEEI LARVVPPDQS FQENYAGIFH FQFWQYGEWV EVVVDRLPT KDGELLFVHS AEGSEFWSAL LEKAYAKING CYEALSGGAT TEGFEDFTGG IAEWYELRKP PPNLFKIIQK ALEKGSLLGC SIDITSAADS EAVTYQKLVK GHAYSVTGAE EVESGSLQK LIRIRNPWGQ VEWTKWVNDN CPSWNTVDPE VRANLTERQE DGEFWMSFSD FLRHYSRLEI CNLTPDTLTC DSYKKWKLTK MDGNWRRGST AGGCRNYPNT FWMNPQYLIK LEEDEDEED GERGCTFLVG LIQKRRRQR KMGEDMHTIG FGIYEVPEEL TGQTNHLGK NFFLTTRARE RSDTFINLRE VLNRFKLPPG EYVLVPSTFE PHKDGDFCIR VFSEKKADYQ AVDDEIEANI EEIDANEEDI DDGFRRLFVQ LAGEDAEISA FELQTILRRV LAKRQDIKSD GFSIETCKIM VDMLDEDGSG KLGLKEFYIL WTKIQKYQKI YREIDVDRSG TMNSYEMRKA LEEAGFKLPC QLHQVIVARF ADDELIIDFD NFVRCLVRLE TLFKIFKQLD PENTGTIQLN LASWLSFSVL
<b>Source</b>	Yeast
<b>Target Names</b>	Capn2
<b>Protein Names</b>	Recommended name: Calpain-2 catalytic subunit EC= 3.4.22.53 Alternative name(s): 80 kDa M-calpain subunit Short name= CALP80 Calcium-activated neutral proteinase 2 Short name= CANP 2 Calpain M-type Calpain-2 large
<b>Expression Region</b>	20-700
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	The calpains, calcium-activated neutral proteases, are nonlysosomal, intracellular cysteine proteases. The mammalian calpains include ubiquitous, stomach-specific, and muscle-specific proteins. The ubiquitous enzymes consist of heterodimers with distinct large, catalytic subunits associated with a common small, regulatory subunit. This gene encodes the large subunit of the ubiquitous enzyme, calpain 2. Multiple heterogeneous transcriptional start sites in the 5' UTR have been reported. Two transcript variants encoding different isoforms



have been found for this gene.

---

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.