



Recombinant Rat Coagulation factor XIII A chain (F13a1)

Product Code	CSB-MP007919RA
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	O08619
Product Type	Recombinant Protein
Immunogen Species	Rattus norvegicus (Rat)
Purity	>85% (SDS-PAGE)
Sequence	GV SLKDYLVNVTAVHLFKERWDS NKIVHHTDKF DNNKLIVRRG QTFYIQIDFN RPYDPRKDLF RVEYVIGRYP QENKGTYPV PVVTELQSGK WGAKVIMNED RSVRLSVQSS PECIVGKFRM YVAVWTPYGI LRTQRDPETD TYILFNPWCE EDAVYLDDEK EREEYVLNDI GGIFHGDFND IKRSWSYGG FEDGILDAFL FVMDKAEMDL SGRGNPIKVS RVGSAMVNAK DDEGVLVGSW DNVYAYGIPP SAWTGSVDIL LEYRSSETPV RYGQCWVLAG VFNTFLRCLG IPARVITNYS SAHDNDANLQ MDIFLEEDGS VSFKLTKDSV WNYHCWNEAW MTRPDLVGVF GGWQAVDSTP QENSDGMYRC GPASVQAVKH GHVCFQFDAP FVLGEVNSDL VYITAKKDGTHVVENVDATH IGKLVITKEI GGDGMQDITD TYKFQEGQEE ERLALETALM YGAKKTLNTE GMVKSSSDVD MNFDVENAVL GKDFRVTITF QNNSNLYTI LAYLSGNITF YTGVSKEFK TESFEVTLDP LSLEKKEVLI RAGEYMSYLL EQGLLHFFVT ARINETRVVL AKQKAIVLTIPKVTIKVRGT AMVGS DMVVT VEFTNPLKET LKNVWLHLEG PGVMRPKRKM FREIRPNATV QWEEVCQPWV SGHRKLNASM TSDSLRHVYG ELDLQIRRRP TV
Source	Mammalian cell
Target Names	F13a1
Protein Names	Recommended name: Coagulation factor XIII A chain Short name= Coagulation factor XIIIa EC= 2.3.2.13 Alternative name(s): Protein-glutamine gamma-glutamyltransferase A chain Transglutaminase A chain
Expression Region	39-732
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	This gene encodes the coagulation factor XIII A subunit. Coagulation factor XIII is the last zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymatic activity and may serve as plasma carrier molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those of plasma origin. Upon cleavage of the activation peptide by thrombin and in the presence



of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin clot. It also crosslinks alpha-2-plasmin inhibitor, or fibronectin, to the alpha chains of fibrin. Factor XIII deficiency is classified into two categories: type I deficiency, characterized by the lack of both the A and B subunits; and type II deficiency, characterized by the lack of the A subunit alone. These defects can result in a lifelong bleeding tendency, defective wound healing, and habitual abortion.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.