



Recombinant Mouse Cytohesin-3 (Cyth3)

Product Code	CSB-MP006469MO
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	O08967
Product Type	Recombinant Protein
Immunogen Species	Mus musculus (Mouse)
Purity	≥85% (SDS-PAGE)
Sequence	MDEGGGGEGG SVPEDLSLEE REELDIRRR KKELIIDIER LKYEIAEVMT EIDNLTVEE SKTTQRNKQI AMGRKKFNMD PPKGIQFLIE NDLLQSSPED VAQFLYKGE G LNKTVIGDY L GERDDFNIV LQAFVELHEF ADLNLVQALR QFLWSFRLPG EAQKIDRMME AFASRYCLCN PGVFQSTDT C YVLSFAIIML NTSLHNHNVR DKPTAERFIT MNRGINEGGD LPEELLRNLY ESIKNEPFKI PEDDGNDLTH TFFNPDREGW LLKLGGRVKT WKRRWFILTD NCLYYFEYTT DKEPRGIPL ENLSIREVED PRKPNCFELY NPSHKGQVIK ACKTEADGRV VEGNHVYRI SAPSPEEKKEE WMKSIKASIS RDPFYDMLAT RKRRRIANKK
Source	Mammalian cell
Target Names	Cyth3
Protein Names	Recommended name: Cytohesin-3 Alternative name(s): ARF nucleotide-binding site opener 3 Short name= Protein ARNO3 General receptor of phosphoinositides 1 Short name= Grp1 PH, SEC7 and coiled-coil domain-containing protein 3
Expression Region	1-399
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	This gene encodes a member of the PSCD (pleckstrin homology, Sec7 and coiled-coil domains) family. PSCD family members have identical structural organization that consists of an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein (GEP) activity, and the PH domain interacts with phospholipids and is responsible for association of PSCDs with membranes. Members of this family appear to mediate the regulation of protein sorting and membrane trafficking. This encoded protein is involved in the control of Golgi structure and function, and it may have a physiological role in regulating ADP-ribosylation factor protein 6 (ARF) functions, in addition to acting on ARF1.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final



concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.