



Recombinant Bovine V-type proton ATPase subunit H (ATP6V1H)

Product Code	CSB-YP002413BO
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	O46563
Product Type	Recombinant Protein
Immunogen Species	Bos taurus (Bovine)
Purity	>85% (SDS-PAGE)
Sequence	<p>MTKMDIRGAV DAAVPTNIIA AKAAEVRANK VNWQSYLQGQ MISSEDCFI QRFEMKRSPE EKQEMLQTEG SQCAKTFINL MTHISKEQTV QYILTLVDDT LQENHQRVSI FFDYAKRSKN TAWSYFLPML NRQDLFTVHM AARIIAKLAA WGKELMEGSD LNYFFNWIKT QLSSQKLRGS GVTAETGTVS SSDSSQYVQC VAGCLQLMLR VNEYRFAWVE ADGVNCIMGV LSNKCGFQLQ YQMIFSVWLL AFSPQMCEHL RRYNIIPVLS DILQESVKEK VTRIILAAFR NFLEKSVERE TRQEYALAMI QCKVLKQLEN LEQQKYDDED ISEDIKFLLE KLGESVQDLS SFDEYSSELK SGRLEWSPVH KSEKFWRENA VRLNEKNYEL LKILTKLLEV SDDPQVLAVA AHDVGEYVRH YPRGKRVIEQ LGGKQLVMNH MHHEDQQVRY NALLAVQKLM VHNWEYLGKQ LQSEQPQTAA ARS</p>
Source	Yeast
Target Names	ATP6V1H
Protein Names	Recommended name: V-type proton ATPase subunit H Short name= V-ATPase subunit H Alternative name(s): V-ATPase 50/57 kDa subunits Vacuolar proton pump subunit H Vacuolar proton pump subunit SFD
Expression Region	1-483
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	<p>This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene encodes the regulatory H subunit of the V1 domain which is required for catalysis of ATP</p>



but not the assembly of V-ATPase. Three alternatively spliced transcript variants encode two isoforms of the H subunit.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.