



# Recombinant Rat Bcl-2-like protein 11 (Bcl2l11)

<b>Product Code</b>	CSB-MP002615RA
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	O88498
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Rattus norvegicus (Rat)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	MAKQPSDVNS ECDREGGQLQ PAERPPQLRP GAP TSLQTES QGNPDGEGDR CPHGSPQGGL APPASGPFA TRSPLFIVR RSSLLSRSS GYFSFDTRS PAMSCDKST QTPSPCQAF NHYLSAMASI RQSQEEPEDL RPEIRIAQEL RRIGDEFNET YTRRAFANDY REAEDHPQMV ILQLLRFIFR LVWRRH
<b>Source</b>	Mammalian cell
<b>Target Names</b>	Bcl2l11
<b>Protein Names</b>	Recommended name: Bcl-2-like protein 11 Short name= Bcl2-L-11 Alternative name(s): Bcl-2-related ovarian death protein Bcl2-interacting mediator of cell death
<b>Expression Region</b>	1-196
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full length protein
<b>Target Details</b>	This protein belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein contains a Bcl-2 homology domain 3 (BH3). It has been shown to interact with other members of the BCL-2 protein family, including BCL2, BCL2L1/BCL-X(L), and MCL1, and to act as an apoptotic activator. The expression of this gene can be induced by nerve growth factor (NGF), as well as by the forkhead transcription factor FKHR-L1, which suggests a role of this gene in neuronal and lymphocyte apoptosis. Transgenic studies of the mouse counterpart suggested that this gene functions as an essential initiator of apoptosis in thymocyte-negative selection. Several alternatively spliced transcript variants of this gene have been identified.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
<b>Shelf Life</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.



Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.