



# Recombinant Mouse Stanniocalcin-2 (Stc2)

<b>Product Code</b>	CSB-MP022822MO
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	O88452
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Mus musculus (Mouse)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	TDSTNP PEGPQDRSSQ QKGRSLQNT AEIQHCLVNA GDVGCVFEC FENNSCEIQG LHGICMTFLH NAGKFDAQGK SFIKDALRCK AHALRHKFGC ISRKCPAIRE MVFQLQRECY LKHDLCSAAQ ENVGIVIVEMI HFKDLLLHEP YVDLVNLLLT CGEDVKEAVT RSVQAQCEQS WGGLCSILSF CTSNIQRPPT AAPEHQPLAD RAQLSRPHHR DTDHHLTANR GAKGERGSKS HPNAHARGRT GGQSAQGPSG SSEWEDEQSE YSDIRR
<b>Source</b>	Mammalian cell
<b>Target Names</b>	Stc2
<b>Protein Names</b>	Recommended name: Stanniocalcin-2 Short name= STC-2
<b>Expression Region</b>	25-296
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	This gene encodes a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. The encoded protein has 10 of its 15 cysteine residues conserved among stanniocalcin family members and is phosphorylated by casein kinase 2 exclusively on its serine residues. Its C-terminus contains a cluster of histidine residues which may interact with metal ions. The protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Constitutive overexpression of human stanniocalcin 2 in mice resulted in pre- and postnatal growth restriction, reduced bone and skeletal muscle growth, and organomegaly. Expression of this gene is induced by estrogen and altered in some breast cancers.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
<b>Shelf Life</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.



Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.