



Recombinant Mouse Signal transducing adapter molecule 1 (Stam)

Product Code	CSB-MP022792MO
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P70297
Product Type	Recombinant Protein
Immunogen Species	Mus musculus (Mouse)
Purity	>85% (SDS-PAGE)
Sequence	<p>PLFATNPFDDQDVEKATSELNNTAEDWGLILDICDKVGGQSRTGPKDCLRSIMRRV NHKDPHV AMQALTLGACVSNCGKIFHLEVCSRDFASEVSNVLNKGHPKVCEKCLKALMVE WTDEFKN DPQLSLISAMIKNLKEQGVTFPAIGSQAAEQAKASPALVAKDPGTVATKKEEED LAKAIE LSLKEQRQQSAPVSTLYPSTSNLLTNHQHEGRKVRAVYDFEAAEDNELTFKAG EIITVLD DSDPNWWKGETHQGVGLFPSNFVTADLTAPEMIKTEKKTQVFNDDVQIETIE PEPEPAF IDEDKMDQLLQMLQSTDPNDQPDPELLHLEAMCHQMGLIDEKLEDIDRKH SELSELN VKVMEALSLYTKLMNEDPMYSMYAKLQSQYYLQSSAVSASQVYPGPAQSG TYLVAGSAQ MTHLQSYSLPPEQLSSISQGAVPSSANQALPSQQTQASYPNAMVSSVQGNYSY PSQASIYS PPAAAAAAAAAAVVPVVPADVTIYQNAGPTMSQVPNYTLTSSTLPQTGGSQ QPPQPQQA YSQKALL</p>
Source	Mammalian cell
Target Names	Stam
Protein Names	Recommended name: Signal transducing adapter molecule 1 Short name= STAM-1
Expression Region	2-548
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	This gene was identified by the rapid tyrosine-phosphorylation of its product in response to cytokine stimulation. The encoded protein contains a SH3 domain and the immunoreceptor tyrosine-based activation motif (ITAM). This protein associates with JAK3 and JAK2 kinases via its ITAM region, and is phosphorylated by the JAK kinases upon cytokine stimulation, which suggests



the function of this protein is as an adaptor molecule involved in the downstream signaling of cytokine receptors. HGS/HRS (hepatocyte growth factor-regulated tyrosine kinase substrate) has been found to bind and counteract the function of this protein.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.