



# Recombinant Human Cadherin-1 (CDH1)

<b>Product Code</b>	CSB-EP005034HU1
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P12830
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	DWVIPP ISCPENEKGP FPKNLVQIKS NKDKKEGKVFY SITGQGADTP PVGVFIIERE TGWLKVTEPL DRERIATYTL FSHAVSSNGN AVEDPMEILI TVTDQNDNKP EFTQEVEFKGS VMEGALPGTS VMEVTATDAD DDVNTYNAAI AYTILSQDPE LPDKNMFTIN RNTGVISVVT TGLDRESFPT YTLVVQAADL QGEGLSSTAT AVITVTDND NPPIFNPTTY KGQVPENEAN VVITTLKVTD ADAPNTPAWE AVYTILNDDG GQFVTTNPV NNDGILKTAK GLDFEAKQQY ILHVAVTNVV PFEVSLTTST ATVTVDVLDV NEAPIFVPPE KRVEVSEDFG VGQEITSYTA QEPDTFMEQK ITYRIWRDTA NWLEINPDTG AISTRAELDR EDFEHVKNST YTALIIATDN GSPVATGTGT LLLILSDVND NAPIPEPRTI FFCERNPKPQ VINIIDADLP PNTSPFTAEL THGASANWTI QYNDPTQESI ILKPKMALEV GDYKINLKLM DNQNKDQVTT LEVSVCDCEG AAGVCRKAQP VEAGLQIPA
<b>Source</b>	E.coli
<b>Target Names</b>	CDH1
<b>Protein Names</b>	Recommended name: Cadherin-1 Alternative name(s): CAM 120/80 Epithelial cadherin Short name= E-cadherin Uvomorulin CD_antigen= CD324 Cleaved into the following 3 chains: 1. E-Cad/CTF1 2. E-Cad/CTF2 3. E-Cad/CTF3
<b>Expression Region</b>	155-709
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Extracellular domain
<b>Target Details</b>	This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function is thought to contribute to progression in cancer by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. Identified transcript variants arise from mutation at consensus splice sites.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a



concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.