



Recombinant Human Calpain-2 catalytic subunit (CAPN2)

Product Code	CSB-YP004496HU
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P17655
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	S HDRAIKYLNQ DYEALRNECL EAGTLFQDPS FPAIPSALGF KELGPYSSKT RGIEWKRPT E ICADPQFIIG GATRTRDQCQ ALGDCWLLAA IASLTLNEEI LARVVPLNQS FQENYAGIFH FQFWQYGEWV EVVVDDRLPT KDGELLFVHS AEGSEFWSAL LEKAYAKING CYEALSGGAT TEGFEDFTGG IAEWYELKKP PPNLFKIIQK ALQKGSLLGC SIDITSAADS EAITFQKLVK GHAYSVTGAE EVESNGSLQK LIRIRNPWGE VEWTRGRWVND CPSWNTIDPE ERERLRRHE DGEFWMFSFD FLRHYSRLEI CNLTPDTLTS DTYKKWKLTK MDGNWRRGST AGGCRNYPNT FWMNPQYLIK LEEDEDEDED GESGCTFLVG LIQKHRRRQR KMGEDMHTIG FGIYEVPEEL SGQTNHLSK NFFLTNRARE RSDTFINLRE VLNRFKLPPG EYILVPSTFE PNKDGDFCIR VFSEKKADYQ AVDDEIEANL EEFDISEDDI DDGFRRLFAQ LAGEDAEISA FELQTLRRV LAKRQDIKSD GFSIETCKIM VDMLDSDGSG KLGLKEFYIL WTKIQKYQKI YREIDVDRSG TMNSYEMRKA LEEAGFKMPC QLHQVIVARF ADDQLIIDFD NFVRCLVRLE TLFKIFKQLD PENTGTIELD LISWLCFSVL
Source	Yeast
Target Names	CAPN2
Protein Names	Recommended name: Calpain-2 catalytic subunit EC= 3.4.22.53 Alternative name(s): Calcium-activated neutral proteinase 2 Short name= CANP 2 Calpain M-type Calpain large polypeptide L2 Calpain-2 large subunit Millimolar
Expression Region	20-700
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	The calpains, calcium-activated neutral proteases, are nonlysosomal, intracellular cysteine proteases. The mammalian calpains include ubiquitous, stomach-specific, and muscle-specific proteins. The ubiquitous enzymes consist of heterodimers with distinct large, catalytic subunits associated with a common small, regulatory subunit. This gene encodes the large subunit of the ubiquitous enzyme, calpain 2. Multiple heterogeneous transcriptional start sites in the 5' UTR have been reported. Two transcript variants encoding different isoforms



have been found for this gene.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.