



Rabbit anti-Mouse IgG Fc Antibody;FITC conjugated

Product Code	CSB-PA00580G0Rb
Abbreviation	Mouse IgG Fc fragment
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Immunogen	Mouse IgG Fc fragment
Raised In	Rabbit
Species Reactivity	Mouse
Relevance	<p>The fragment crystallizable region (Fc region) is the tail region of an antibody that interacts with cell surface receptors called Fc receptors and some proteins of the complement system. This property allows antibodies to activate the immune system. In IgG, IgA and IgD antibody isotypes, the Fc region is composed of two identical protein fragments, derived from the second and third constant domains of the antibody's two heavy chains; IgM and IgE Fc regions contain three heavy chain constant domains (CH domains 2–4) in each polypeptide chain. The Fc regions of IgGs bear a highly conserved N-glycosylation site. Glycosylation of the Fc fragment is essential for Fc receptor-mediated activity. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and α-2,6 linked sialic acid residues. The other part of an antibody, called the Fab region, contains variable sections that define the specific target that the antibody can bind. By contrast, the Fc region of all antibodies in a class are the same for each species; they are constant rather than variable. The Fc region is, therefore, sometimes incorrectly termed the "fragment constant region".</p>
Form	Liquid
Conjugate	FITC conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Caprylic Acid Ammonium Sulfate Precipitation purified
Isotype	IgG
Clonality	Polyclonal
Product Type	Secondary Antibody