



# Recombinant Human Histone deacetylase 2 (HDAC2)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-BP010238HU  |
| <b>Abbreviation</b>      | HDAC2   |
| <b>Storage</b>           | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.<br>Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.  |
| <b>Uniprot No.</b>       | Q92769  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Homo sapiens (Human)  |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | MAYSQGGGKK KVCYYYDGDI GNYYYYGQGHP MKPHRIRMTH NLLLNYGLYR<br>KMEIYRPHKA TAEEMTKYHS DEYIKFLRSI RPDNMSEYSK QMQRFNVGED<br>CPVFDGLFEF CQLSTGGSVA GAVKLNRRQT DMAVNWAGGL<br>HHAKKSEASG FCYVNDIVLA ILELLKYHQR VLYIDIDIHH GDGVVEAFYT<br>TDRVMTVSFH KYGEYFPGTG DLRDIGAGKG KYYAVNFPMPR DGIDDESYGQ<br>IFKPIISKVM EMYQPSAVVL QCGADSLSGD RLGCFLNLTVK GHAKCVEVVK<br>TFNLPLLMLG GGGYTIRNVA RCWTYETAVA LDCEIPNELP YNDYFEYFGP<br>DFKLHISPSN MTNQNTPEYM EKIKQRLFEN LRMLPHAPGV QMQAIPEDAV<br>HEDSGDEDGE DPKKRISIRA SDKRIACDEE FSDSEDEGEG GRRNVADHKK<br>GAKKARIEED KKETEDKKT D VKEEDKSKDN SGEKTDTKGT KSEQLSNP |
| <b>Source</b>            | Baculovirus   |
| <b>Target Names</b>      | HDAC2   |
| <b>Protein Names</b>     | Recommended name: Histone deacetylase 2 Short name= HD2 EC= 3.5.1.98  |
| <b>Expression Region</b> | 1-488   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | full length protein   |
| <b>Target Details</b>    | This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes and are responsible for the deacetylation of lysine residues on the N-terminal region of the core histones (H2A, H2B, H3 and H4). This protein also forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus it plays an important role in transcriptional regulation, cell cycle progression and developmental events.                         |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a  |



concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.