



# Recombinant Human CD2-associated protein (CD2AP)

<b>Product Code</b>	CSB-YP004914HU
<b>Abbreviation</b>	CD2AP
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	Q9Y5K6
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	≥85% (SDS-PAGE)
<b>Sequence</b>	MVDYIVEYDY DAVHDELTI RVGEIIRNVK KLQEEGWLEG ELNGRRGMFP DNFVKEIKRE TEFKDDSLPI KRERHGNVAS LVQRISTYGL PAGGIQPHPQ TKNIKKTKK RQCKVLF EYI PQNEDELELK VGDIIDINEE VEEGWWSGTL NNLGLFPSN FVKELEVTDG GETHEAQDDS ETVLAGPTSP IPSLGNVSET ASGSVTQPKK IRGIGFGDIF KEGSVKLRTR TSSSETEKK PEKPLILQSL GPKTQSVEIT KTDTEGKIKI KEYCRTLFAY EGTNEDELTF KEGEIIHLIS KETGEAGWWR GELNGKEGVF PDNFAVQINE LDKDFPKPKK PPPPAKAPAP KPELIAAEKK YFSLKPEEKD EKSTLEQKPS KPAAPQVPPK KPTPPTKASN LLRSSGTVYP KRPEKPVPPP PPIAKINGEV SSISSKFETE PVSKLKL DSE QLPLRPKSVD FDSLTVRTSK ETDVVNFDDI ASSENLLHLT ANRPKMPGRR LPGRFNGGHS PTHSPEKILK LPKEEDSANL KPSELKKDTC YSPKPSVYLS TPSSASKANT TAFLTPLEIK AKVETDDVKK NSLDELRAQI IELLCIVEAL KKD HGKELEK LRKDL EEEKT MRSNLEMEIE KLK KAVLSS
<b>Source</b>	Yeast
<b>Target Names</b>	CD2AP
<b>Protein Names</b>	Recommended name: CD2-associated protein Alternative name(s): Adapter protein CMS Cas ligand with multiple SH3 domains
<b>Expression Region</b>	1-639
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	full length protein
<b>Target Details</b>	This gene encodes a scaffolding molecule that regulates the actin cytoskeleton. The protein directly interacts with filamentous actin and a variety of cell membrane proteins through multiple actin binding sites, SH3 domains, and a proline-rich region containing binding sites for SH3 domains. The cytoplasmic protein localizes to membrane ruffles, lipid rafts, and the leading edges of cells.



It is implicated in dynamic actin remodeling and membrane trafficking that occurs during receptor endocytosis and cytokinesis. Haploinsufficiency of this gene is implicated in susceptibility to glomerular disease.

---

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.