



# Recombinant Human Galactoside 3 (4)-L-fucosyltransferase (FUT3), partial

<b>Product Code</b>	CSB-EP009077HU1
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P21217
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	RVSRDDATGSPRAPSGSSRQDTPTRPTLLILLWTWPFHIPVALSRCSEMVPG TADCHIT ADRKVYPQADTVIVHHWDIMSNPKSRLPPSPRPQGQRWIWFNLEPPPNCQHL EALDRYFN LTMSYRSDSDIFTPYGWLEPWGQPAHPLNLSAKTELVAWAVSNWKPDSAR VRYYQSLQ AHLKVDVYGRSHKPLPKGTMETLSRYKFYLAFENSLHPDYITEKLWRNALEA WAVPVVL GPSRSNYERFLPPDAFIHVDDFQSPKDLARYLQELDKDHARYLSYFRWRETLR PRFSWA LDFCKACWKLQQESRYQTVRSIAAWFT
<b>Source</b>	E.coli
<b>Target Names</b>	FUT3
<b>Protein Names</b>	Recommended name: Galactoside 3(4)-L-fucosyltransferase EC= 2.4.1.65 Alternative name(s): Blood group Lewis alpha-4-fucosyltransferase Short name= Lewis FT Fucosyltransferase 3 Fucosyltransferase III Short name= FucT-I
<b>Expression Region</b>	35-361
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	partial
<b>Target Details</b>	The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metastasis, inflammation, and bacterial adhesion. They are secondarily absorbed to red blood cells giving rise to their Lewis phenotype. This gene is a member of the fucosyltransferase family, which catalyzes the addition of fucose to precursor polysaccharides in the last step of Lewis antigen biosynthesis. It encodes an enzyme with alpha(1,3)-fucosyltransferase and alpha(1,4)-fucosyltransferase activities. Mutations in this gene are responsible for the majority of Lewis antigen-negative phenotypes. Multiple alternatively



spliced variants, encoding the same protein, have been found for this gene.

---

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.