



Recombinant Human 26S protease regulatory subunit 6A (PSMC3)

Product Code	CSB-YP018891HU
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P17980
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	MNLLPNIESP VTRQEKMATV WDEAEQDGIG EEVLKMSTEE IIQRTRLLDS EIKIMKSEVL RVTHELQAMK DKIKENSEKI KVNKTLPLYLV SNVIELLDVD PNDQEEDGAN IDLDSQRKKGK CAVIKTSTRQ TYFLPVIGLV DAEKCLKPGDL VGVNKDSYLI LETLPTEYDS RVKAMEVDER PTEQYSDIGG LDKQIQELVE AIVLPMNHKE KFENLGIQPP KGVLMYGPFG TGKTLLARAC AAQTKATFLK LAGPQLVQMF IGDGAKLVRD AFALAKEKAP SIIFIDELDA IGTKRFDSEK AGDREVQRTM LELLNQLDGF QPNTQVKVIA ATNRVDILDP ALLRSGRLDR KIEFPMPEE ARARIMQIHS RKMNVSPDVN YEELARCTDD FNGAQCKAVC VEAGMIALRR GATELTHEDY MEGILEVQAK KKANLQYYA
Source	Yeast
Target Names	PSMC3
Protein Names	Recommended name: 26S protease regulatory subunit 6A Alternative name(s): 26S proteasome AAA-ATPase subunit RPT5 Proteasome 26S subunit ATPase 3 Proteasome subunit P50 Tat-binding protein 1 Short name= TBP-1
Expression Region	1-439
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the



viral protein and the transcription complex. A pseudogene has been identified on chromosome 9.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.