



Recombinant Pig Interleukin-12 subunit alpha (IL12A)

| | |
|--------------------------|--|
| Product Code | CSB-YP641818PI |
| Abbreviation | IL12A |
| Storage | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. |
| Uniprot No. | Q29053 |
| Product Type | Recombinant Protein |
| Immunogen Species | Sus scrofa (Pig) |
| Purity | ≥85% (SDS-PAGE) |
| Sequence | RSLPA TTAGPGMFKC LNHSQNLLKA VSNTLQKAKQ TLEFYSCTSE EIDHEDITKD KTSTVEACL P LELATNESCL AARETSLITN GNCLTSGKTS FMTTLCLSSI YEDLKMYHVE FQAMNAKLLM DPKRQIFLDQ NMLTAITELM QALNFNSETV P QKPSLEELD FYKTKIKLCI LLHAFRIRAV TIDRMMSYLN SS |
| Source | Yeast |
| Target Names | IL12A |
| Protein Names | Recommended name: Interleukin-12 subunit alpha Short name= IL-12A Alternative name(s): Cytotoxic lymphocyte maturation factor 35 kDa subunit Short name= CLMF p35 IL-12 subunit p35 |
| Expression Region | 26-222 |
| Notes | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| Tag Info | Tag type will be determined during the manufacturing process. |
| Protein Length | Full Length of Mature Protein |
| Target Details | This gene encodes a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final |



concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.