



Recombinant Bovine Myc proto-oncogene protein (MYC)

Product Code	CSB-EP645319BO-B
Abbreviation	MYC
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q2HJ27
Product Type	Recombinant Protein
Immunogen Species	Bos taurus (Bovine)
Purity	>85% (SDS-PAGE)
Sequence	MPLNVSFANK NYDLDYDSVQ PYFYCDEEEN FYHQQQQSEL QPPAPSEDIW KKFELLPTTP LSPSRRSGLC SPSYVAVASF SPRGDDDGGG GSFSSADQLE MVTELLGGDM VNQSFICDPD DETLIKNIII QDCMWSGFSA AAKLVSEKLA SYQAARKDGG SPSPARGHGG CSTSSLYLQD LSAAASECID PSVVFYPLN DSSSPKPCAS PDSTAFSPSS DSSLSSAESS PRASPEPLAL HEETPPTTSS DSEEEQEDEE EIDVVSVEKR QPPAKRSESG SPSAGSHSKP PHSPLVLKRC HVSTHQHNYA APPSTRKDYP AAKRAKLD SG RVLKQISNNR KCASPRSSDT EENDKRRTHN VLERQRRNEL KRSFFALRDQ IPELENNEKA PKVVILKKAT AYILSVQAEQ QCLKSEIDVL QKRREQLKLK LEQIRNSCA
Source	E.coli
Target Names	MYC
Protein Names	Recommended name: Myc proto-oncogene protein Alternative name(s): Proto-oncogene c-Myc Transcription factor p64
Expression Region	1-439
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	full length protein
Target Details	This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in



Burkitt's lymphomas, suggesting its importance in the normal function of this gene.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.