



Recombinant Mouse MAP kinase-activated protein kinase 3 (Mapkapk3)

Product Code	CSB-EP659737MO
Abbreviation	Mapkapk3
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q3UMW7
Product Type	Recombinant Protein
Immunogen Species	Mus musculus (Mouse)
Purity	>85% (SDS-PAGE)
Sequence	MDGETAGEKG SLVPPPGALG GSALGGAPAP GVRREPKKYA VTDDYQLSKQ VLGLGVNGKV LECYHRRSGQ KCALKLLYDS PKARQEVDDH WQASGGPHIV RILDVYENMH HGKRCLLIVM ECEGGELFS RIQERGDQAF TEREAAEIMR DIGTAIQFLH SRNIAHRDVK PENLLYTSKE KDAVLKLTDF GFAKETTQNA LQTPCYTPYY VAPEVLGPEK YDKSCDMWSL GVIMYILLCG FPPFYSNTGQ AISPGMKRRI RLGQYSFPNP EWLDVSEDAK QLIRLLLKTD PTERLTIMQF MNHPWINQSM VVPQTPLYTA RVLQEDKDHV DDKVEEMTSA LATMRVDYDQ VKIKDLKTSN NLLLNKRRKK QAGSSSASQG CNNQ
Source	E.coli
Target Names	Mapkapk3
Protein Names	Recommended name: MAP kinase-activated protein kinase 3 Short name= MAPK-activated protein kinase 3 Short name= MAPKAP kinase 3 Short name= MAPKAP-K3 Short name= MAPKAPK-3 Short name= MK-3 EC= 2.7.11.1
Expression Region	1-384
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	full length protein
Target Details	This gene encodes a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)- activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and



repress the activity of E47, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and cell differentiation.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.