



Recombinant Bovine Cyclin-H (CCNH)

| | |
|--------------------------|---|
| Product Code | CSB-EP670250BO-B |
| Abbreviation | CCNH |
| Storage | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. |
| Uniprot No. | Q3ZBL9 |
| Product Type | Recombinant Protein |
| Immunogen Species | Bos taurus (Bovine) |
| Purity | ≥85% (SDS-PAGE) |
| Sequence | MYHNSSQKRH WTFASEEQLA RLRADANRKF KCKAVANGKV LPNDPVFLEP HEEITLCKYY EKRLLEFCSV FKPAMPRSVV GTACMYFKRF YLNNSVMEYH PRIIMLTCAF LACKVDEFNV SSPQFVGNLR ESPLGQEKTL EQILEYELL IQLNFHLIV HNPYRPFEGF LIDLKTRYPL LENPEILRKT ADDFLNRVAL TDAHLLYTPS QIALTAILSS ASRAGITMES YLSESLMLKE NRTSLSQLLD IMKSMRNLVK KYEPPRPEEV AALKQKLERC HSAELALNVV TKKRKGYEDD DYVSKKSKHE EVCSPKGSFM |
| Source | E.coli |
| Target Names | CCNH |
| Protein Names | Recommended name: Cyclin-H |
| Expression Region | 1-320 |
| Notes | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| Tag Info | Tag type will be determined during the manufacturing process. |
| Protein Length | full length protein |
| Target Details | This protein belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final |



concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.