



Recombinant Rat 26S protease regulatory subunit 7 (Psmc2)

Product Code	CSB-YP720404RA
Abbreviation	Psmc2
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q63347
Product Type	Recombinant Protein
Immunogen Species	Rattus norvegicus (Rat)
Purity	≥85% (SDS-PAGE)
Sequence	PDYLGADQR KTKEDEKDDK PIRALDEGDI ALLKTYGQST YSRQIKQVED DIQQLLKKIN ELTGIKESDT GLAPPALWDL AADKQTLQSE QPLQVARCTK IINADSEDPK YIINVKQFAK FVVDLSDQVA PTDIEEGMRV GVDRNKYQIH IPLPPKIDPT VTMMQVEEKP DVTYSVGGC KEQIEKLREV VETPLLHPER FVNLGIEPPK GVLLFGPPGT GKTLCARAVA NRTDACFIPV IGSELVQKYV GEGARMVREL FEMARTKKAC LIFFDEIDAI GGARFDDGAG GDNEVQRTML ELINQLDGFDP RGNIKVLMA TNRPDTLDPALMRPGRDRK IEFSLPDLEG RTHIFKIHAR SMSVERDIRF ELLARLCPNS TGAEIRSVCT EAGMFAIRAR RKIATEKDFL EAVNKVIKSY AKFSATPRYM TYN
Source	Yeast
Target Names	Psmc2
Protein Names	Recommended name: 26S protease regulatory subunit 7 Alternative name(s): 26S proteasome AAA-ATPase subunit RPT1 Proteasome 26S subunit ATPase 2 Protein MSS1
Expression Region	2-433
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal



pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit has been shown to interact with several of the basal transcription factors so, in addition to participation in proteasome functions, this subunit may participate in the regulation of transcription. This subunit may also compete with PSMC3 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.