



Recombinant Human 60S ribosomal protein L22 (RPL22)

Product Code	CSB-YP020182HU
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P35268
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	APVKKLVVK GGKKKKQVLK FTLDCSTEPVE DGIMDAANFE QFLQERIKVN GKAGNLGGGV VTIERSKSKI TVTSEVPFSK RYLKYLTKKY LKKNNLRDWL RVVANSKESY ELRYFQINQD EEEEEDED
Source	Yeast
Target Names	RPL22
Protein Names	Recommended name: 60S ribosomal protein L22 Alternative name(s): EBER-associated protein Short name= EAP Epstein-Barr virus small RNA-associated protein Heparin-binding protein HBp15
Expression Region	2-128
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22E family of ribosomal proteins. Its initiating methionine residue is post-translationally removed. The protein can bind specifically to Epstein-Barr virus-encoded RNAs (EBERs) 1 and 2. The mouse protein has been shown to be capable of binding to heparin. Transcript variants utilizing alternative polyA signals exist. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. It was previously thought that this gene mapped to 3q26 and that it was fused to the acute myeloid leukemia 1 (AML1) gene located at 21q22 in some therapy-related myelodysplastic syndrome patients with 3;21 translocations; however, these fusions actually involve a ribosomal protein L22 pseudogene located at 3q26, and this gene actually maps to 1p36.3-p36.2.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a



concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.