



# Recombinant Human Prostaglandin G/H synthase 2 (PTGS2), partial

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-BP018986HU  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P35354  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Homo sapiens (Human)  |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | ANPCCSHPCQNRGVCMSVGFQYKDCDTRTGFYGENCSTPEFLTRIKLFLKP<br>TPNTVHYILTHFKGFVNWNIPFLRNAIMSYVLTSRSHLIDSPPTYNADYGYK<br>SWEAFSNLSYYTRALPPVPDDCPTPLGVKGGKQLPDSNEIVEKLLLRKFIPDP<br>QGSNMMFAFFAQHFTHQFFKTDHKRGAFTNGLGHGVDLNHIYGETLARQRK<br>LRLFKDGMKYQIIDGEMYPPTVKDTQAEMIPPQVPEHLRFVAVGQEVFGLVP<br>GLMMYATIWLREHNRVCDVLKQEHPEWGDEQLFQTSRLILIGETIKIVIEDYVQ<br>HLSGYHFKLKFDPELLFNKQFQYQNRIAAEFNTLYHWHPLLPDTFQIHDQKYN<br>YQQFIYNNILLSILEHGITQFVESFTRQIAGR VAGGRNVPPAVQKVSQASIDQSRQ<br>MKYQSFNEYRKRFLKPYESFEELTGEKEMSAELEALYGDIDAVELYPALLVE<br>KPRPDAIFGETMVEVGAPFSLKGLMGNVICSPAYWKPSTFGGEVGFQIINTASI<br>QSLICNNVKGCPFTSFSVPDPELIKTVTINASSRSGLDDINPTVLLKERS |
| <b>Source</b>            | Baculovirus   |
| <b>Target Names</b>      | PTGS2   |
| <b>Protein Names</b>     | Recommended name: Prostaglandin G/H synthase 2 EC= 1.14.99.1<br>Alternative name(s): Cyclooxygenase-2 Short name= COX-2 PHS II Prostaglandin H2<br>synthase 2 Short name= PGH synthase 2 Short name= PGHS-2 Prostagla   |
| <b>Expression Region</b> | 18-601  |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Partial   |
| <b>Target Details</b>    | Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis.  |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final   |



concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.