



# Recombinant Mouse Tyrosine-protein phosphatase non-receptor type 13 (Ptpn13), partial

|                          |  |
|--------------------------|--|
| <b>Product Code</b>      | CSB-MP734108MO   |
| <b>Abbreviation</b>      | Ptpn13   |
| <b>Storage</b>           | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.<br>Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.   |
| <b>Uniprot No.</b>       | Q64512   |
| <b>Product Type</b>      | Recombinant Protein  |
| <b>Immunogen Species</b> | Mus musculus (Mouse)   |
| <b>Purity</b>            | >85% (SDS-PAGE)  |
| <b>Source</b>            | Mammalian cell   |
| <b>Target Names</b>      | Ptpn13   |
| <b>Protein Names</b>     | Recommended name: Tyrosine-protein phosphatase non-receptor type 13 EC= 3.1.3.48 Alternative name(s): PTP36 Protein tyrosine phosphatase DPZPTP Protein tyrosine phosphatase PTP-BL Protein-tyrosine phosphatase RIP   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.  |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.  |
| <b>Protein Length</b>    | Partial  |
| <b>Target Details</b>    | This protein is a member of the protein tyrosine phosphatase (PTP) family. PTPs are signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP is a large intracellular protein. It has a catalytic PTP domain at its C-terminus and two major structural domains: a region with five PDZ domains and a FERM domain that binds to plasma membrane and cytoskeletal elements. This PTP was found to interact with, and dephosphorylate, Fas receptor and I $\kappa$ B $\alpha$ through the PDZ domains. This suggests it has a role in Fas mediated programmed cell death. This PTP was also shown to interact with GTPase-activating protein, and thus may function as a regulator of Rho signaling pathways. Four alternatively spliced transcript variants, which encode distinct proteins, have been reported. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.  |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients,  |



storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.