



# Recombinant Sheep Isocitrate dehydrogenase [NADP] cytoplasmic (IDH1)

<b>Product Code</b>	CSB-BP751079SH
<b>Abbreviation</b>	IDH1
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	Q6XUZ5
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Ovis aries (Sheep)
<b>Purity</b>	≥85% (SDS-PAGE)
<b>Sequence</b>	SHKIQGGSV VEMQGDDEMTR IIWELIKEKL IFPYVDLDLH SYDLSIENRD ATNDQVTKDA AEAIKKYNVG VKCATITPDE KRVEEFKLLKQ MWKSPNGTIR NILGGTVFRE AIICKNIPRL VSGWVKPIII GRHAYGDQYR ATDFVVPGPG KVEICYTPSD GSPKTVYL VH N FTESGGVAM GMFNQDKSIE DFAHSSFQMA LSKNWPLYLS TKNTILKKYD GRFKDIFQEI YDKQYKSQFE AQNIWYEHRL IDDMVAQAMK SEGGFIWACK NYDGDVQSDS VAQGYGSLGM MTSVLVCPDG KTVEAEAAHG TVTRHYRMYQ KGQETSTNPI ASIFAWTRGL AHRAKLDNNK ELSFFAKALE EVCIETIEAG FMTKDLAACI KGLPNVQRSD YLNTFEFMDK LGENLQLKLA QAKL
<b>Source</b>	Baculovirus
<b>Target Names</b>	IDH1
<b>Protein Names</b>	Recommended name: Isocitrate dehydrogenase [NADP] cytoplasmic Short name= IDH EC= 1.1.1.42 Alternative name(s): Cytosolic NADP-isocitrate dehydrogenase IDP NADP(+)-specific ICDH Oxalosuccinate decarboxylase
<b>Expression Region</b>	2-414
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. This protein is the NADP(+)-dependent isocitrate dehydrogenase



found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production.

---

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.