



Recombinant Pig Proteasome activator complex subunit 2 (PSME2)

Product Code	CSB-BP768115PI
Abbreviation	PSME2
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q863Z0
Product Type	Recombinant Protein
Immunogen Species	Sus scrofa (Pig)
Purity	≥85% (SDS-PAGE)
Sequence	AKPCGVRLS GEARKQVDVF RQNLFQEAE FLYRFLPQKI IYLSQLLQED SLNVTDLTSL RAPLDIIPD PPPKDDMET DKQEKKVVPK CGFLPGNEKI LALLGLVKPE VWTLKEKCIL VITWQHLP KIEDGNDFGV AVQEKVLERV NAVKTKEAF QTTISKYFSE RGDAAKASK ETHVMDYRAL VHERDEAAHG ELRAMVLDLR AFYAELYHII SSNLEKIVDP KGEEKPSMY
Source	Baculovirus
Target Names	PSME2
Protein Names	Recommended name: Proteasome activator complex subunit 2 Alternative name(s): Proteasome activator 28 subunit beta Short name= PA28b Short name= PA28beta
Expression Region	2-239
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta



and gamma) of the 11S regulator have been identified. This gene encodes the beta subunit of the 11S regulator, one of the two 11S subunits that is induced by gamma-interferon. Three beta and three alpha subunits combine to form a heterohexameric ring. Six pseudogenes have been identified on chromosomes 4, 5, 8, 10 and 13.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.