



# Recombinant Rabbit Stromelysin-1 (MMP3)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-YP014676RB  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P28863  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Oryctolagus cuniculus (Rabbit)  |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | FSTFPGTPKW TKTHLTYRIV NYTPDLPRDA VDAAIEKALK VWEVTPPLTF<br>SRKYEGEADI MISFGVREHG DFIPFDGPGN VLAHAYAPGP GINGDAHFD<br>DEQWTKDTTG TNLFLVAAHE LGHSLGLFHS ANPEALMYPV YNAFTDLARF<br>RLSQDDVDGI QSLYGPAPAS PDNSGVPMEP VPPGSGTPVM CDPDLSFDAI<br>STLRGEILFF KDRYFWRKSL RILEPEFHLL SSWWPSLPSA VDAAYEVISR<br>DTVFIFKGTQ FWAIRGNEVQ AGYPRSIHTL GFPSTIRKID AAISDKERKK<br>TYFFVEDKYW RFDEKRQSLE PGFPRHIAED FPGINPKIDA VFEAFGFFYF<br>FSGSSQSEFD PNAKKVTHVL KSNSWFQC   |
| <b>Source</b>            | Yeast   |
| <b>Target Names</b>      | MMP3  |
| <b>Protein Names</b>     | Recommended name: Stromelysin-1 Short name= SL-1 EC= 3.4.24.17<br>Alternative name(s): Matrix metalloproteinase-3 Short name= MMP-3 Transin-1   |
| <b>Expression Region</b> | 101-478   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Full Length of Mature Protein   |
| <b>Target Details</b>    | Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP s are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients,   |



storage temperature and the stability of the protein itself.  
Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.