



# Recombinant Rat Coagulation factor VII (F7)

<b>Product Code</b>	CSB-EP811708RA-B
<b>Abbreviation</b>	F7
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	Q8K3U6
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Rattus norvegicus (Rat)
<b>Purity</b>	≥85% (SDS-PAGE)
<b>Sequence</b>	ANSLLEELW SSSLERECNE ERCSFEEARE IFKSPERTKQ FWTIYSDGDQ CASNPCQNGG TCQDHLKSYV CFCPLDFEGR NCEKNKNEQL ICANENGDCD QYCRDHVGTK RTCSCHEDYV LQPDEVSCKP KVEYPCGRIP VVEKRNF SRP QGR
<b>Source</b>	E.coli
<b>Target Names</b>	F7
<b>Protein Names</b>	Recommended name: Coagulation factor VII EC= 3.4.21.21 Alternative name(s): Serum prothrombin conversion accelerator Cleaved into the following 2 chains: 1. Factor VII light chain 2. Factor VII heavy chain
<b>Expression Region</b>	42-193
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	This gene encodes coagulation factor VII which is a vitamin K-dependent factor essential for hemostasis. This factor circulates in the blood in a zymogen form, and is converted to an active form by either factor IXa, factor Xa, factor XIIa, or thrombin by minor proteolysis. Upon activation of the factor VII, a heavy chain containing a catalytic domain and a light chain containing 2 EGF-like domains are generated, and two chains are held together by a disulfide bond. In the presence of factor III and calcium ions, the activated factor then further activates the coagulation cascade by converting factor IX to factor IXa and/or factor X to factor Xa. Alternative splicing of this gene results in 2 transcripts. Defects in this gene can cause coagulopathy.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final



concentration of glycerol is 50%. Customers could use it as reference.

### Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.