



Recombinant Human NAD-dependent protein deacetylase sirtuin-2 (SIRT2)

Product Code	CSB-EP812889HU
Abbreviation	SIRT2
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q8IXJ6
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	≥85% (SDS-PAGE)
Sequence	AEPDPSHPL ETQAGKVQEA QDSDSDSEGG AAGGEADMDF LRNLFSQTLS LGSQKERLLD ELTLEGVARY MQSERCRRVI CLVGAGISTS AGIPDFRSPS TGLYDNLEKY HLPYPEAIFE ISYFKKHPEP FFALAKELYP GQFKPTICHY FMRLKDKGL LLRCYTQNIID TLERIAGLEQ EDLVEAHGTF YTSHCVSASC RHEYPLSWMK EKIFSEVTPK CEDCQSLVKP DIVFFGESLP ARFFSCMQSD FLKVDLLLVM GTSLQVQPFA SLISKAPLST PRLINKEKA GQSDPFLGMI MGLGGGMDFD SKKAYRDVAW LGECDQGCLA LAELLGWKKE LEDLVRREHA SIDAQSGAGV PNPSTSASPK KSPPPAKDEA RTTEREKPQ
Source	E.coli
Target Names	SIRT2
Protein Names	Recommended name: NAD-dependent protein deacetylase sirtuin-2 EC= 3.5.1.- Alternative name(s): Regulatory protein SIR2 homolog 2 SIR2-like protein 2
Expression Region	2-389
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. This protein is included in class I of the sirtuin family. Two transcript variants result from alternative splicing of this gene.

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.