



# Recombinant Chicken Fibroblast growth factor 8 (FGF8)

<b>Product Code</b>	CSB-YP845867CH
<b>Abbreviation</b>	FGF8
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	Q90722
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Gallus gallus (Chicken)
<b>Purity</b>	≥85% (SDS-PAGE)
<b>Sequence</b>	QVTVQSPP NFTQHVREQS LVTDQLSRRL VRTYQLYSRT SGKHVQILDN KKINAMAEDG DVHAKLIVET DTFGSRVRIK GAATGFYICM NKKGKLGKGS NGKGKDCVFT EIVLENNYTA LQNAKYEGWY MAFTRKGRPR KGSKTRQHQR EVHFMKRLPK GHQTTEPHRR FEFLNYPFNR RSKRTRNSSA SLRP
<b>Source</b>	Yeast
<b>Target Names</b>	FGF8
<b>Protein Names</b>	Recommended name: Fibroblast growth factor 8 Short name= FGF-8 Alternative name(s): Heparin-binding growth factor 8 Short name= HBGF-8
<b>Expression Region</b>	23-214
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	This protein is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogenesis. The adult expression of this gene is restricted to testes and ovaries. Temporal and spatial pattern of this gene expression suggests its function as an embryonic epithelial factor. Studies of the mouse and chick homologs revealed roles in midbrain and limb development, organogenesis, embryo gastrulation and left-right axis determination. The alternative splicing of this gene results in four transcript variants.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a



concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

### Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.