



Recombinant Human DNA damage-binding protein 2 (DDB2)

Product Code	CSB-BP846067HU
Abbreviation	DDB2
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q92466
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	MAPKKRPETQ KTSEIVLRPR NKRSRSPLEL EPEAKKLC AK GSGPSRRCDSDCLWVGLAGP QILPPCRSIV RTLHQHKLGR ASWPSVQQGL QQSFLHTLDS YRILQKAAPF DRATSLAWH PTHPSTVAVG SKGGDIMLWN FGIKDKPTFI KGIGAGGSIT GLKFNPLNTN QFYASSMEGT TRLQDFKGN I LRVFASSDTI NIWFCSLDVS ASSRMVVTGD NVGNVILLNM DGKELWNL RM HKKKVTHVAL NPCCDWFLAT ASVDQTVKIW DLRQVRGKAS FLYSLPHRHP VNAACFSPDG ARLLTTDQKS EIRVYSASQW DCPLGLIPHP HRHFQHLTPI KAAWHPRYNL IVVGRYPDPN FKSCPTYELR TIDVFDGNSG KMMCQLYDPE SSGISSLNEF NPMGDTLASA MGYHILIWSQ EEARTRK
Source	Baculovirus
Target Names	DDB2
Protein Names	Recommended name: DNA damage-binding protein 2 Alternative name(s): DDB p48 subunit Short name= DDBb Damage-specific DNA-binding protein 2 UV-damaged DNA-binding protein 2 Short name= UV-DDB 2
Expression Region	1-427
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	full length protein
Target Details	This gene encodes a protein that is necessary for the repair of ultraviolet light-damaged DNA. This protein is the smaller subunit of a heterodimeric protein complex that participates in nucleotide excision repair, and this complex mediates the ubiquitylation of histones H3 and H4, which facilitates the cellular response to DNA damage. This subunit appears to be required for DNA binding. Mutations in this gene cause xeroderma pigmentosum complementation group E, a recessive disease that is characterized by an increased sensitivity to UV light and a high predisposition for skin cancer development, in some cases



accompanied by neurological abnormalities.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.