



Recombinant Human E3 ubiquitin-protein ligase TRIM63 (TRIM63)

| | |
|--------------------------|--|
| Product Code | CSB-MP846575HU |
| Abbreviation | TRIM63 |
| Storage | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. |
| Uniprot No. | Q969Q1 |
| Product Type | Recombinant Protein |
| Immunogen Species | Homo sapiens (Human) |
| Purity | ≥85% (SDS-PAGE) |
| Sequence | MDYKSSLIQD GNP MENLEKQ LICPICLEMF TKPVVILPCQ HNL CRKCAND IFQAANPYWT SRGSSVMSG GRFRCPTCRH EVIMDRHGVY GLQRNLLVEN IIDYKQEC S RPLQKGSHP MCKEHEDEKI NIYCLTCEVP TCSMCKVFGI HKACEVAPLQ SVFQGQKTEL NNCISMLVAG NDRVQTIITQ LEDSRRTKE NSHQVKEELS QKFDTLYAIL DEKSELLQR ITQE QEKLS FIALIQQYQ EQLDKSTKLV ETAIQLDEP GGATFLLTAK QLIKSIVEAS KGCQLGKTEQ GFENMDFFTL DLEHIADALR AIDFGTDEEE EEFIEEEDQE EEESTEGKEE GHQ |
| Source | Mammalian cell |
| Target Names | TRIM63 |
| Protein Names | Recommended name: E3 ubiquitin-protein ligase TRIM63 EC= 6.3.2.- Alternative name(s): Iris RING finger protein Muscle-specific RING finger protein 1 Short name= MuRF-1 Short name= MuRF1 RING finger protein 28 Stria |
| Expression Region | 1-353 |
| Notes | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| Tag Info | Tag type will be determined during the manufacturing process. |
| Protein Length | full length protein |
| Target Details | This gene encodes a member of the RING zinc finger protein family found in striated muscle and iris. The product of this gene is localized to the Z-line and M-line lattices of myofibrils, where titin s N-terminal and C-terminal regions respectively bind to the sarcomere. In vitro binding studies have shown that this protein also binds directly to titin near the region of titin containing kinase activity. Another member of this protein family binds to microtubules. Since these family members can form heterodimers, this suggests that these proteins may serve as a link between titin kinase and microtubule-dependent signal pathways in muscle. |



Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.