



# Recombinant Mouse Macrophage metalloelastase (Mmp12)

<b>Product Code</b>	CSB-MP014659MO
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P34960
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Mus musculus (Mouse)
<b>Purity</b>	≥85% (SDS-PAGE)
<b>Sequence</b>	R SRWMKRYLTY RIYNYTPDMK REDVDYIFQK AFQVWSDVTP LRFRKLHKDE ADIMILFAFG AHGDFNYFDG KGGTLAHAFY PGPGIQGDAH FDEAETWTKS FQGTNLFLVA VHELGHSLGL QHSNNPKSIM YPTYRYLNPS TFRLSADDIR NIQSLYGAPV KPPSLTKPSS PPSTFCHQSL SFDAVTTVGE KIFFFKDWFF WWKLPGSPAT NITSISSIWP SIPSGIQAAY EIESRNQLFL FKDEKYWLIN NLVPEPHYPR SIYSLGFSAS VKKVDAAVFD PLRQKVYFFV DKHYWRYDVR QELMDPAYPK LISTHFPGIK PKIDAVLYFK RHYYIFQGAY QLEYDPLFRR VTKTLKSTSW FGC
<b>Source</b>	Mammalian cell
<b>Target Names</b>	Mmp12
<b>Protein Names</b>	Recommended name: Macrophage metalloelastase Short name= MME EC= 3.4.24.65 Alternative name(s): Matrix metalloproteinase-12 Short name= MMP-12
<b>Expression Region</b>	110-473
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP s are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. It is thought that This protein is cleaved at both ends to yield the active enzyme, but this processing has not been fully described. The enzyme degrades soluble and insoluble elastin. It may play a role in aneurysm formation and studies in mice suggest a role in the development of emphysema. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final



concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

---

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.