





Recombinant Human Testican-3 (SPOCK3)

Product Code	CSB-BP859412HU
Abbreviation	SPOCK3
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q9BQ16
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	AAAAVAAAG GRSDGGNFLD DKQWLTTISQ YDKEVGQWNK FRDEVEDDYF RTWSPGKPFD QALDPAKDPC LKMKCSRHKV CIAQDSQTAV CISHRRLTHR MKEAGVDHRQ WRGPILSTCK QCPVVYPSPV CGSDGHTYSF QCKLEYQACV LGKQISVKCE GHCPCPSDKP TSTSRNVKRA CSDLEFREVA NRLRDWFKAL HESGSQNKKT KTLLRPERSR FDTSILPICK DSLGWMFNRL DTNYDLLLDQ SELRSIYLDK NEQCTKAFFN SCDTYKDSLI SNNEWCYCFQ RQQDPPCQTE LSNIQKRQGV KKLLGQYIPL CDEDGYYKPT QCHGSVGQCW CVDRYGNEVM GSRINGVADC AIDFEISGDF ASGDFHEWTD DEDDEDDIMN DEDEIEDDDE DEGDDDDGGD DHDVYI
Source	Baculovirus
Source Target Names	Baculovirus SPOCK3
Target Names	SPOCK3 Recommended name: Testican-3 Alternative name(s): SPARC/osteonectin,
Target Names Protein Names	SPOCK3 Recommended name: Testican-3 Alternative name(s): SPARC/osteonectin, CWCV, and Kazal-like domains proteoglycan 3
Target Names Protein Names Expression Region	SPOCK3 Recommended name: Testican-3 Alternative name(s): SPARC/osteonectin, CWCV, and Kazal-like domains proteoglycan 3 22-436 Repeated freezing and thawing is not recommended. Store working aliquots at
Target Names Protein Names Expression Region Notes	SPOCK3 Recommended name: Testican-3 Alternative name(s): SPARC/osteonectin, CWCV, and Kazal-like domains proteoglycan 3 22-436 Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Target Names Protein Names Expression Region Notes Tag Info	SPOCK3 Recommended name: Testican-3 Alternative name(s): SPARC/osteonectin, CWCV, and Kazal-like domains proteoglycan 3 22-436 Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag type will be determined during the manufacturing process.