



Recombinant Dog Acyl-CoA-binding protein (DBI)

| | |
|--------------------------|--|
| Product Code | CSB-EP866124DO |
| Abbreviation | DBI |
| Storage | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. |
| Uniprot No. | Q9TQX6 |
| Product Type | Recombinant Protein |
| Immunogen Species | Canis lupus familiaris (Dog) (Canis familiaris) |
| Purity | >85% (SDS-PAGE) |
| Sequence | SQAEFDKAA EDVKHLKTKP ADDEMLYIYS HYKQATVGDI NTERPGLLDL RGKAKWDAWN QLKGTSKEDA MKAYVNKVED LKKKYGI |
| Source | E.coli |
| Target Names | DBI |
| Protein Names | Recommended name: Acyl-CoA-binding protein Short name= ACBP Alternative name(s): Diazepam-binding inhibitor Short name= DBI Endozepine Short name= EP |
| Expression Region | 2-87 |
| Notes | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| Tag Info | Tag type will be determined during the manufacturing process. |
| Protein Length | Full Length of Mature Protein |
| Target Details | This gene encodes diazepam binding inhibitor, a protein that is regulated by hormones and is involved in lipid metabolism and the displacement of beta-carbolines and benzodiazepines, which modulate signal transduction at type A gamma-aminobutyric acid receptors located in brain synapses. The protein is conserved from yeast to mammals, with the most highly conserved domain consisting of seven contiguous residues that constitute the hydrophobic binding site for medium- and long-chain acyl-Coenzyme A esters. Diazepam binding inhibitor is also known to mediate the feedback regulation of pancreatic secretion and the postprandial release of cholecystokinin, in addition to its role as a mediator in corticotropin-dependent adrenal steroidogenesis. Three pseudogenes located on chromosomes 6, 8 and 16 have been identified. Multiple transcript variants encoding different isoforms have been described for this gene. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final |



concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.