



# Recombinant Human Regulator of nonsense transcripts 2 (UPF2), partial

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-BP867186HU  |
| <b>Abbreviation</b>      | UPF2  |
| <b>Storage</b>           | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.<br>Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.  |
| <b>Uniprot No.</b>       | Q9HAU5  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Homo sapiens (Human)  |
| <b>Purity</b>            | ≥85% (SDS-PAGE)   |
| <b>Source</b>            | Baculovirus   |
| <b>Target Names</b>      | UPF2  |
| <b>Protein Names</b>     | Recommended name: Regulator of nonsense transcripts 2 Alternative name(s): Nonsense mRNA reducing factor 2 Up-frameshift suppressor 2 homolog Short name= hUpf2   |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Partial   |
| <b>Target Details</b>    | This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located in the perinuclear area. It interacts with translation release factors and the proteins that are functional homologs of yeast Upf1p and Upf3p. Two splice variants have been found for this gene; both variants encode the same protein. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.<br>Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.  |