



Recombinant Chicken Dual specificity protein phosphatase 4 (DUSP4)

Product Code	CSB-EP879388CH
Abbreviation	DUSP4
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q9PW71
Product Type	Recombinant Protein
Immunogen Species	Gallus gallus (Chicken)
Purity	>85% (SDS-PAGE)
Sequence	MVAPEGLREM EGSALRRLVG REEASGGRCL LLDCRPFLAH SAGHIRGALN VRCNTIVRRR AKGAVSLEQI LPAEGEVRRAR LRAGLYTAVV LYDERSPRAE ALRDDSTVAL VLRALRRDMA RADIRLLAGG YERFASEYPE FCAKTKTLSS ISPPSSAESL DLGFSSCGTP LHDQGGPVEI LPFLYLGSAY HAARRDMLDA LGITALLNVS SDCPNHFEGH YQYKCIPVED NHKADISSWF MEAIEYIDSV KECCGRVLVH CQAGISRSAT ICLAYLMMKK RVKLEKAFEF VKQRRSIISP NFSFMGQLLQ FESQVLATSC AVEAASPSGT LRER GKATST PTSQVFVFSFP VSVG VHATPS SLPYLHSPIT TSPSC
Source	E.coli
Target Names	DUSP4
Protein Names	Recommended name: Dual specificity protein phosphatase 4 EC= 3.1.3.16 EC= 3.1.3.48 Alternative name(s): Mitogen-activated protein kinase phosphatase 2 Short name= MAP kinase phosphatase 2 Short name= MKP-2
Expression Region	1-375
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	full length protein
Target Details	This protein is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product



inactivates ERK1, ERK2 and JNK, is expressed in a variety of tissues, and is localized in the nucleus. Two alternatively spliced transcript variants, encoding distinct isoforms, have been observed for this gene. In addition, multiple polyadenylation sites have been reported.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.