







Recombinant Human Von Hippel-Lindau disease tumor suppressor (VHL)

Product Code CSB-MP025852HU Storage Store at -20°C, for extended storage, conserve at -20°C or -80°C. Uniprot No. P40337 Product Type Recombinant Protein Immunogen Species Homo sapiens (Human) Purity >85% (SDS-PAGE) Sequence MPRRAENWDE AEVGAEEAGV EEYGPEEDGG EESGAEESGP EESGAPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPQPYPT LPPGTGRRIH SYRGHLWLFR DAGTHDGLLV NQTELFVPSL NYDGQPIFAN ITLPVYTLKE RCLQVVRSLV KPENYRRLDI VRSLYEDLED HPNVQKDLER LTQERIAHQR MGD Source Mammalian cell Target Names VHL Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullfin-2, and possesses ubiquitin ligase E3 activity. This protein is involved i		
Uniprot No. P40337 Product Type Recombinant Protein Immunogen Species Homo sapiens (Human) Purity >85% (SDS-PAGE) Sequence MPRRAENWDE AEVGAEEAGV EEYGPEEDGG EESGAEESGP EESGPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPDGPQPYPT LPPGTGRRIH SYRGHLWIFR DAGTHDGLLV NQTELFVPSL NVDGQPIFAN ITLPVYTLKE RCLQVVRSLV KPENYRRLDI VRSLYEDLED HPNVQKDLER LTQERIAHQR MGD Source Mammalian cell Target Names VHL Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working allquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POL	Product Code	CSB-MP025852HU
Product Type Recombinant Protein Immunogen Species Homo sapiens (Human) Purity >85% (SDS-PAGE) Sequence MPRRAENWDE AEVGAEEAGV EEYGPEEDGG EESGAEESGP EESGPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPCPYPT LPPGTCRIH SYRGHLWLFR DAGTHDGLLV NOTELFVPSL NYDGQPIPAN ITLPVYTLER RCLQVVRSLV KPENYRRLDI VRSLYEDLED HPNVQKDLER LTQERIAHQR MGD Source Mammalian cell Target Names VHL Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPBT is also reported to be a target of this protein. Alternatively spli	Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Immunogen Species	Uniprot No.	P40337
Purity >85% (SDS-PAGE) Sequence MPRRAENWDE AEVGAEEAGV EEYGPEEDGG EESGAEESGP EESGPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPQPYPT LPPGTGRRIH SYRGHLWUFR DAGTHDGLLV NQTELFVPSL NVDGQPIFAN ITLPVYTLKE RCLQVVRSLV KPENYRRLDI VRSLYEDLED HPNVQKDLER LTQERIAHQR MGD Source Mammalian cell Target Names VHL Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration) and aliquot for long-term storage at -20°C-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Product Type	Recombinant Protein
Sequence MPRRAENWDE AEVGAEEAGV EEYGPEEDGG EESGAEESGP EESGPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPQPYPT LPPGTGRRIH SYRGHLWUFR DAGTHDGLLV NQTELFVPSL NVDGQPIFAN ITLPVYTLKE RCLQVVRSLV KPENYRRLDI VRSLYEDLED HPNVQKDLER LTQERIAHQR MGD Source Mammalian cell Target Names VHL Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please re	Immunogen Species	Homo sapiens (Human)
EESGPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPQPYPT LPPGTGRRIH SYRGHLWLFR DAGTHDGLLV NQTELFVPSL NVDGQPIFAN ITLPVYTLKE RCLQVVRSLV KPENYRRLDI VRSLYEDLED HPNVQKDLER LTQERIAHQR MGD Source Mammalian cell Target Names VHL Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Purity	>85% (SDS-PAGE)
Target Names VHL Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Sequence	EESGPEELGA EEEMEAGRPR PVLRSVNSRE PSQVIFCNRS PRVVLPVWLN FDGEPQPYPT LPPGTGRRIH SYRGHLWLFR DAGTHDGLLV NQTELFVPSL NVDGQPIFAN ITLPVYTLKE RCLQVVRSLV KPENYRRLDI VRSLYEDLED
Protein Names Recommended name: Von Hippel-Lindau disease tumor suppressor Alternative name(s): Protein G7 pVHL Expression Region 1-213 Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Source	Mammalian cell
Notes Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.	Target Names	VHL
Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Protein Names	··
Tag Info Tag type will be determined during the manufacturing process. Protein Length Full length protein Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Expression Region	1-213
Protein Length Full length protein Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Notes	
Target Details Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Tag Info	Tag type will be determined during the manufacturing process.
syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Protein Length	Full length protein
contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference. Shelf Life The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.	Target Details	syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritance of VHL syndrome. This protein is a component of the protein complex that includes elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB7 is also reported to be a target of this protein. Alternatively spliced transcript variants
storage temperature and the stability of the protein itself.	Reconstitution	contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final
	Shelf Life	



CUSABIO TECHNOLOGY LLC





Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.